

**COMPOUND INORGANIC MICRONUTRIENT FERTILISER: [PFC 1(C)(II)(b)]
MINERAL MICRONUTRIENT FERTILISER IN SOLUTION**

DECLARED CONTENT:

4.0 % w/wWater-soluble Boron (B) from boron ethanalamine
0.15 % w/wWater-soluble Copper (Cu), from sulphate
0.015 % w/wWater-soluble Molybdenum (Mo), from sodium
4.5 % w/wWater-soluble Zinc (Zn), from citrate

Unintentional N and P₂O₅ content:

4.0% w/wNitrogen (N)

LIST OF INGREDIENTS:

Boron ethanalamine* (CAS no 68425-67-2), Zinc citrate* (CAS no. 546-46-3),
*Virgin material substances and mixtures.

INSTRUCTIONS FOR USE:

SugarMover Zn is a liquid fertiliser that provides boron, copper, molybdenum and zinc, the main nutrients required in the grain and bulb filling process. The boron present in **SugarMover Zn** is an essential element that improves the transport of sugars from the leaves (sources) to the grains and bulbs (sinks). Copper, molybdenum and zinc are nutrients that promote photosynthesis and protein synthesis that improves quality, caliber, and uniformity characteristics. Thanks to **Stoller Formulation Technology**, we provide adequate nutrition that naturally intervenes in the physiological processes of plants.

INSTRUCTIONS AND APPLICATION RATES							
Crop					Growth stage		
Cereals	Wheat, barley, rye, triticale, oats...	✓	0.65 %	300-500	2-3 L/ha	From seed filling up to maturity (BBCH 71-89)	1-2 applications every 15-30 days
	Sorghum	✓	0.65 %	300-500	2-3 L/ha	From seed filling up to maturity (BBCH 71-89)	1-2 applications every 15-30 days
	Corn	✓	0.65 %	300-500	2-3 L/ha	From seed filling up to maturity (BBCH 71-89)	1-2 applications every 15-30 days
	Sunflower seed	✓	0.65 %	300-500	2-3 L/ha	From seed filling up to maturity (BBCH 71-89)	1-2 applications every 15-30 days
Oilseed crops and pulses	Soybean	✓	0.30 %	700-1000	2-3 L/ha	From pods appearance to maturity (BBCH 71-89)	1-2 applications every 15-30 days
	Rapeseed	✓	0.30 %	700-1000	2-3 L/ha	From pods appearance to maturity (BBCH 71-89)	1-2 applications every 15-30 days
Grasses	Pea, bean	✓	0.65 %	300-500	2-3 L/ha	10 days before cutting	1 application
	Lucerne	✓	0.65 %	300-500	2-3 L/ha	10 days before cutting	1 application
Tubers	✓	0.30 %	700-1000	2-3 L/ha	From tuber filling up to maturity (BBCH 71-89)	1-2 applications every 15-20 days	
Bulbs	✓	0.30 %	700-1000	2-3 L/ha	From bulb filling up to maturity (BBCH 71-89)	1-2 applications every 15-20 days	
Root vegetables	✓	0.30 %	700-1000	2-3 L/ha	From root filling up to maturity (BBCH 71-89)	1-2 applications every 15-20 days	
Foliar application	Irrigation application	Seed treatment	Furrow treatment				
Optimal dose	Dosage min - max	Treatment volume (L)	Treat. Vol (L) min - max	Frequency			

If you have any questions, please contact Stoller Europe, or the distributor of Stoller Europe, or your technical advisor, for more specific recommendations.

Observations:

Carefully read the instructions on the label before using the product. The product is compatible with the most common fertilisers. We recommend to previously consult with your distributor or local representative. To be used only where there is a recognized need. Do not exceed the application rate. These doses are indicative, and it will be the user's responsibility to increase or decrease them according to the crop's characteristics, always following the advice of a specialist technician. In any case, we recommend making a small-scale test before using the product.

RECOMMENDED STORAGE CONDITIONS:

Store according to local legislation. Observe indications on the label. Store the containers between 0 and 35 °C, in a dry and well-ventilated place, far from sources of heat and direct solar light. Keep far away from ignition points. Keep away from oxidizing agents and from highly acidic or alkaline materials. Do not smoke. Prevent the entry of non-authorized persons. Once the containers are open, they must be carefully closed and placed vertically to prevent spills. Please also refer to the information provided in the Material Safety Data Sheet (MSDS).

SAFETY AND ENVIRONMENTAL INFORMATION:

Carefully read the instructions on the label before using the product. To avoid risks to human health and the environment, comply with the instructions for use. Labelling in accordance with Regulation (EC) No 1272/2008:



ANGER

Hazard statements:
H315. Causes skin irritation.
H335. May cause respiratory irritation.
H411. Toxic to aquatic life with long lasting effects.

Precautionary statements:
P264. Wash zones in contact thoroughly after handling.
P273. Avoid release to the environment.
P280. Wear protective gloves/protective clothing/eye protection/face protection/hearing protection/...
P303+P361+P353. IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water [or shower].
P305+P351+P338. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

OTHER INFORMATION/ADDITIONAL INFORMATION:

Professional use.
Fertiliser usable in organic plant production according to R(EU) 2018/848.

WARNING:

STOLLER EUROPE S.L.U. makes all the recommendations based on experience. A lot of uncontrollable factors (preparation of mixtures, climatology, application forms, etc.) may be involved. Therefore, the user will be responsible for any damage that may occur.

Stoller
Together we grow.

Manufactured by:
STOLLER EUROPE, S.L.U.
A Stoller Group Company
Elche Parque Empresarial
C/ Max Planck, 1.
03203 Elche, Alicante (Spain)
(+34) 965 110 522
info@stollereurope.com
www.stollereurope.com



This document is a reference tool and may only be used as such. The appropriate product label is the one shown on the packaging and printed at the time of purchase.

PHYSICAL FORM: Liquid in solution
NET MASS: 26.2 kg
NET VOLUME: 20 L
LOT NUMBER:

