

**STRAIGHT INORGANIC MICRONUTRIENT FERTILISER: [PFC 1(C)(II)(a)]  
MICRONUTRIENT SOLUTION FERTILISER**

**DECLARED CONTENT:**

6.0% w/w ..... Water-soluble Molybdenum (Mo), from sodium

**Unintentional N and P<sub>2</sub>O<sub>5</sub> content:**

3.6% w/w ..... Nitrogen (N)

**LIST OF INGREDIENTS:**

Sodium Molybdate<sup>1</sup> (CAS no. 10102-40-6), Monoethanolamine<sup>1</sup> (CAS no. 141-43-5)

<sup>1</sup>Virgin material substances and mixtures.

**INSTRUCTIONS FOR USE:**

**Keylate Mo** is a liquid fertiliser developed to counteract molybdenum deficiencies. Molybdenum is a fundamental element for the absorption of nitrogen in plants. The **Stoller Technology** present in the formulation allows the product to be absorbed, translocated and assimilated in a more efficient way.

INSTRUCTIONS AND APPLICATION RATES							
Crop	Application Method	Concentration	Volume	Rate	Growth stage	Frequency	
Fruiting vegetables	Protected	✓	0.43%	700-1000	3-5 L/ha	From sprouting to harvest	5-12 Applications every 7-15 days 2-6 Applications every 15-30 days
	Open field	✓					
Leaf vegetables	Protected	✓	0.43%	700-1000	3-5 L/ha	From sprouting to harvest	5-12 Applications every 7-15 days 2-6 Applications every 15-30 days
	Open field	✓					
Tubers	✓	0.43%	700-1000	3-5 L/ha	From sprouting to harvest	2-6 Applications every 15-30 days	
Bulbs	✓	0.43%	700-1000	3-5 L/ha	From sprouting to harvest		
Root vegetables	✓	0.43%	700-1000	3-5 L/ha	From sprouting to harvest		
Orchards	Stone fruit	✓	0.30%	1000-1500	3-5 L/ha	From sprouting to harvest	2-6 Applications every 15-30 days
	Pome fruit	✓					
	Tropical	✓					
	Nuts	✓					
	Persimmon	✓	0.15%	2000-2500			
	Pomegranate	✓					
Sapling trees	✓	0.30%	700-1000	2-3 L/ha	From sprouting to harvest	2-6 Applications every 15-30 days	
Citrus	✓	0.12%	2500-3000	3-5 L/ha	From sprouting to harvest	2-6 Applications every 15-30 days	
Olive trees	✓	0.15%	2000-2500	3-5 L/ha	From sprouting to harvest		
Grapes	Table grapes	✓	0.15%	2000-2500	3-5 L/ha	From sprouting to harvest	2-6 Applications every 15-30 days
	Vines	✓	0.60%	500-700			
Berries	Protected	✓	0.43%	700-1000	3-5 L/ha	From sprouting to harvest	2-6 Applications every 7-15 days
	Open field	✓					
Strawberry	Protected	✓	0.43%	700-1000	3-5 L/ha	From sprouting to harvest	
	Open field	✓					

- Foliar application
- Irrigation application
- Seed treatment
- Furrow treatment
- Optimal dose
- Dosage min - max
- Treatment volume (L)
- Treat. Vol (L) min - max
- Frequency

If you have any questions, please contact Stoller Europe, or the distributor of Stoller Europe, or your technical advisor, for more specific recommendations.

**Observations:**

Carefully read the instructions on the label before using the product.

**Keylate Mo** is designed to be used through foliar applications. The product is compatible with most liquid fertilisers.

However, if mixing is desired, it is advisable to consult your local distributor or representative in advance.

Use only in cases of known necessity. Do not exceed the application rate. These doses are merely indicative, and it will be the user's responsibility to increase or decrease them according to the crop's characteristics, always following the advice of a specialist technician. In any case, we recommend making a small-scale test before using the product.

**RECOMMENDED STORAGE CONDITIONS:**

Store according to local legislation. Observe indications on the label. Store the containers between 0 and 35 °C, in a dry and well-ventilated place, far from sources of heat and direct solar light. Keep far away from ignition points. Keep away from oxidising agents and from highly acidic or alkaline materials. Do not smoke. Prevent the entry of non-authorised persons. Once the containers are open, they must be carefully closed and placed vertically to prevent spills. Please also refer to the information provided in the Material Safety Data Sheet (MSDS).

**SAFETY AND ENVIRONMENTAL INFORMATION:**

**Carefully read the instructions on the label before using the product.** To avoid risks to human health and the environment, comply with the instructions for use.

Labelling in accordance with Regulation (EC) No 1272/2008:



**DANGER**

**PRODUCT CLASSIFICATION:**

Hazard statements:

- H315. Causes skin irritation.
- H318. Causes serious eye damage.
- H335. May cause respiratory irritation.

Precautionary statements:

- P280. Wear protective gloves/protective clothing/eye protection/face protection/hearing protection/...
- P305+P351+P338. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
- P310. Immediately call a POISON CENTER/doctor if you feel unwell.
- P362+P364. Take off contaminated clothing and wash it before reuse.
- P403+P233. Store in a well-ventilated place. Keep container tightly closed.
- P501. Dispose of contents/container to a chemical waste bin.

**OTHER INFORMATION/ADDITIONAL INFORMATION:**

Professional use.

**WARNING:**

STOLLER EUROPE S.L.U. makes all the recommendations based on experience. A lot of uncontrollable factors (preparation of mixtures, climatology, application forms, etc.) may be involved. Therefore, the user will be responsible for any damage that may occur.



Manufactured by:  
**STOLLER EUROPE, S.L.U.**  
A Stoller Group Company  
Elche Parque Empresarial  
C/ Max Planck, 1.  
03203 Elche, Alicante (Spain)  
(+34) 965 110 522  
info@stollereurope.com  
www.stollereurope.com

**This document is a reference tool and may only be used as such. The appropriate product label is the one shown on the packaging and printed at the time of purchase.**



**PHYSICAL FORM:** Liquid in solution  
**NET MASS:** 5.8 kg  
**NET VOLUME:** 5 L  
**LOT NUMBER:**