

**COMPOUND LIQUID INORGANIC MACRONUTRIENT FERTILISER: [PFC 1 (C)(I)(b)(ii)]
MINERAL FERTILISER NK 10.5-5 WITH MICRONUTRIENTS IN SOLUTION**

DECLARED CONTENT:

10.5 % w/wTotal nitrogen (N)
10.5 % w/wUreic nitrogen (N)
5.0 % w/wWater-soluble potassium oxide (K ₂ O)
0.002 % w/wWater-soluble cobalt (Co) from EDTA
0.002 % w/wWater-soluble molybdenum (Mo) from sodium

LIST OF INGREDIENTS:

Urea¹ (CAS: 57-13-6), Potassium Citrate¹(CAS: 6100-05-6).
¹Virgin material substances and mixture.

INSTRUCTIONS FOR USE:

BioForge is a binary liquid fertiliser that can be used as a source of both nitrogen and potassium, as well as the microelements cobalt and molybdenum, essential nutrients for optimal plant development and to favor its resistance in difficult periods. The **Stoller Technology** present in the formulation of the product provides adequate nutrition to intervene naturally in the physiological processes of the plants.

INSTRUCTIONS AND APPLICATION RATES						
Crop						Frequency
Arables Cereals, corn, oil seed crops	✓	0.10 %	300-500	0.3-0.5	In mixture with herbicides. During unfavorable periods.	1-2 applications every 30-60 days
Fruiting vegetables	Protected	✓	0.08 %	700-1000	0.5-1	1-6 applications every 7-15 days
	Open field	✓	0.15 %	700-1000	1-1.5	Weekly 1-3 applications every 15-30 days
Leaf vegetables	Protected	✓	0.08 %	700-1000	0.5-1	1-6 applications every 7-15 days
	Open field	✓	0.15 %	700-1000	1-1.5	Weekly 1-3 applications every 30-60 days
Root vegetables, tubers, bulbs	✓	0.15 %	700-1000	1-1.5	In mixture with herbicides. During unfavorable periods.	1-3 applications every 15-30 days
Pulses	✓	0.20 %	250-500	0.5-1	During growth cycle. During unfavorable periods.	1-3 applications every 15-30 days
Orchards Stone fruit, pome fruit Persimmon, pomegranate, tropical nuts	✓	0.05 %	2000-2500	1-1.5	During growth cycle. During unfavorable periods.	1-4 applications every 15-30 days
Sapling trees	✓	0.15 %	300-1000	0.5-1.5	Transplanting and during all crop cycle. During unfavorable periods.	1-4 applications every 15-30 days
Citrus	✓	0.04 %	2500-3000			
Olive trees	✓	0.05 %	2000-2500	1-1.5	During growth cycle. During unfavorable periods.	
Grapes Table grapes Vines	✓	0.15 %	700-1000			
	✓	0.20 %	500-700			
Banana	✓	1 L/ha		1-2	Before fruit color change	
Berries, strawberry	✓	0.15 %	700-1000	1-1.5	Transplanting and during growth cycle from first flowering. During unfavorable periods.	1-6 applications every 7-15 days

If you have any questions, please contact Stoller Europe, or the distributor of Stoller Europe, or your technical advisor, for more specific recommendations.

Observations:

The product is compatible with most liquid fertilisers. However, if mixing is desired, it is advisable to consult your local distributor or representative in advance. These doses are indicative, and it will be the user's responsibility to increase or decrease them according to the crop's characteristics, always following the advice of a specialist technician. In any case, we recommend making a small-scale test before using the product. To be used only where there is a recognised need. Do not exceed the application rate.

RECOMMENDED STORAGE CONDITIONS:

Store the containers between 0 and 35 ° C. General conditions for storage: avoid sources of heat, radiation, static electricity and contact with food.
Please also refer to information provided in Material Safety Data Sheet (MSDS).

SAFETY AND ENVIRONMENTAL INFORMATION:

Carefully read the instructions on the label before using the product. To avoid risks to human health and the environment, comply with the instructions for use.
Classification of this product has been carried out in accordance with CLP Regulation (EC) No 1272/2008.



WARNING

PRODUCT CLASSIFICATION:

Hazard statements: H302+H312+H332: Harmful if swallowed, in contact with skin or if inhaled. H319: Causes serious eye irritation. **Precautionary statements:** P264: Wash hands thoroughly after handling. P301+P312: IF SWALLOWED: Call a POISON CENTER/doctor if you feel unwell. P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P312: Call a POISON CENTER/doctor if you feel unwell. P337+P313: If eye irritation persists: Get medical advice/attention. P501: Dispose of contents/ container in accordance with local/regional/national/international regulation.

This fertiliser contains urea, which can release ammonia and have an impact on air quality. Depending on local conditions, appropriate remediation measures must be taken.

OTHER INFORMATION/ADDITIONAL INFORMATION:

Professional use.

Warning:

Stoller Europe S.L.U. makes all the recommendations based on experience. A lot of uncontrollable factors (preparation of mixtures, climatology, application forms, etc.) may be involved. Therefore, the user will be responsible for any damage that may occur.

This document is a reference tool and may only be used as such. The appropriate product label is the one shown on the packaging and printed at the time of purchase.

Manufactured by:
STOLLER EUROPE, S.L.U.
A Stoller Group Company

Elche Parque Empresarial
C/ Max Planch, 1.
03203 Elche, Alicante (Spain)
(+34) 965 110 522
info@stollereurope.com
www.stollereurope.com



PHYSICAL FORM:	Liquid in solution
NET MASS:	5.8 kg e
NET VOLUME:	5 L e
LOT NUMBER:	