

# STRAIGHT LIQUID INORGANIC MACRONUTRIENT FERTILISER: [PFC 1 (C)(I)(b)(i)] MINERAL FERTILISER SOLUTION N ( $C_{a}$ -Mg) 10 (7-2) WITH MICRONUTRIENTS

### DECLARED CONTENT:

10.0% w/w	
10.0% w/w	Ureic nitrogen (N)
	Water-soluble calcium oxide (CaO)
2.0% w/w	Water-soluble magnesium oxide (MgO)
0.1% w/w	
1.0% w/w	Water-soluble zinc (7n) from gluconate

#### LIST OF INGREDIENTS:

Urea¹ (CAS: 57-13-6), Calcium Gluconate¹ (CAS: 299-28-5), Magnesium Gluconate¹ (CAS: 3632-91-5) 
¹Virgin material substances and mixtures.

#### INSTRUCTIONS FOR USE:

**BudPower** is a liquid fertiliser composed of macro and micronutrients which are necessary for a homogeneous and quality flowering. **BudPower** is specially formulated for use just after dormancy in fruit trees, vines and vegetables. The **Stoller Technology** present in the formulation of the product provides adequate nutrition to intervene naturally in the physiological processes of the plants.

	INIC	3.31	CTIO	IC AND	ΑD	NI ICATIONI BATES		
INSTRUCTIONS AND APPLICATION RATES								
	Crop	%		ŧΙ		Growth stage		
Arables	Corn, Sunflower					BBCH vegetative development stages		
Oil seed crops	Rapeseed	~	1.00 %	300-500	3-5	14-18. BBCH Flowering Stages 51-69 (R1-R3)		
Orchards	Stone fruit, pome fruit, persimmon, pomegranate, kiwi, tropical, nuts	~	0.70 %	700-1000	5-10	From swollen leaf bud to swollen flower buds BBCH 03-51		
Sapling trees	3	~	0.60 %	500-700	3-5	From swollen leaf bud to developed bud BBCH 03-39	1-2 applications every 7-15 days	
Citrus		~	0.35 %	1500-2000				
Grapes	Table grapes	~	1.70 %	300-500	_ 5-10	From swollen leaf bud to swollen flower buds BBCH 03-51		
	Vines	~						
Olive trees		~		0.70 % 700-1000				
Berries		~	0.70 %			From vegetative development to the first inflorescences		
Foliar application O Irrigation application		Seed treatment     Furrow treatment						
Optimal dose Dosage min - max (L/ha)		Treatment volume (L)		me (L) Treat. Vol (L) min - max	Frequency			

If you have any questions, please contact Stoller Europe, or the distributor of Stoller Europe, or your technical advisor, for more specific recommendations.

#### Observation

BudPower can be applied to pruning wood before the anticipated bud break, 2 - 4 weeks before bud break until the beginning of flowering or on vegetables before bloom. The product is compatible with most liquid fertiliser, in nutritioning program applications, the mixture should be made in the following order: 1st Water, 2st BudPower and 3st Agrochemicals. Any mixture should be made and applied at the same time and we recommend to previously, consult with your distributor or, local representative. To be used only where there is a recognised need. Do not exceed the application rate. These does are indicative, and it will be the user's responsibility to increase or decrease them according to the crop's characteristics, always following the advice of a specialist technician. In any case, we recommend making a small-scale test before using the product.

#### RECOMMENDED STORAGE CONDITIONS:

Store the containers between 0 and 35 °C. General conditions for storage: avoid sources of heat, radiation, static electricity and contact with food.

Please also refer to information provided in Material Safety Data Sheet (MSDS).

#### SAFETY AND ENVIRONMENTAL INFORMATION:

Carefully read the instructions on the label before using the product. To avoid risks to human health and the environment, comply with the instructions for use.

Classification of this product has been carried out in accordance with CLP Regulation (EC) No 1272/2008.



# Boric Acid (CAS: 10043-35-3)

#### PRODUCT CLASSIFICATION:

Hazard statements: H412: Harmful to aquatic life with long lasting effects. H319: Causes serious eye irritation. H360FD: May damage fertility. May damage free unborn oildi H315: Causes skin irritation. Precautionary statements: P201: Obtain special instructions before use. P280: Wear protective gloves/face protection/protective dothing/respiratory protection/protective footwear. P302+P352: IF ON SKIN: Wash with plenty of water. P303+P351+P338: IF IN EYES. Rines cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P308+P331: IF exposed or concerned: Get medical advice/attention. P501: Dispose of contents/container in accordance with regulations on hazardous waster or packaging and packaging water respectively.

This fertiliser contains urea, which can release ammonia and have an impact on air quality. Depending on local conditions, appropriate remediation measures must be taken.

# OTHER INFORMATION/ADDITIONAL INFORMATION:

Professional use.

# Warning:

Stoller Europe S.L.U. makes all the recommendations based on experience. A lot of uncontrollable factors (preparation of mixtures, climatology, application forms, etc.) may be involved. Therefore, the user will be responsible for any damage that may occur.

# Together we grow.



Manufactured by: STOLLER EUROPE, S.L.U. A Stoller Group Company Elche Parque Empresarial C/ Max Planck, 1. 03203 Elche, Alicante (Spain) (+34) 965 110 522 info@stollereurope.com www.stollereurope.com This document is a reference tool and may only be used as such. The appropiate product label is the one shown on the packaging and printed at the time of purchase.



PHYSICAL FORM: Liquid in solution

NET MASS: 6.5 kg ⊕

51 e

NET VOLUME: LOT NUMBER: