

STRAIGHT LIQUID INORGANIC MACRONUTRIENT FERTILISER: [PFC 1 (C)(I)(b)(i)] MINERAL FERTILISER N (Ca) 15 (12) IN SOLUTION

DECLARED CONTENT:

 15.0 % w/w
 ...Nitrogen (N)

 15.0 % w/w
 ...Ureic nitrogen (N)

 12.0 % w/w
 ...Water-soluble calcium oxide (CaO)

LIST OF INGREDIENTS:

Urea1 (CAS: 57-13-6), Calcium Gluconate1 (CAS: 299-28-5).

¹Virgin material substances and mixture.

INSTRUCTIONS FOR USE:

Harbest is a liquid fertiliser that provides nitrogen and calcium in cereals and extensive crops. These two nutrients ensure the nutritional contribution necessary for the correct development of the crop. Thanks to Stoller N-HIBS patented technology, the urea nitrogen present in Harbest is controlled release making its absorption more efficient.

	INS	ΤR	UCTIO	ONS AND	APPLI	CATION RATES	
Crop		9		Ē		Growth stage	
Arables	Wheat, triticale, barley, rye	~	2.50 %	200-400		From tillering (BBCH 14) to maturity (BBCH 79)	
	Corn, sunflower					From first 3-4 leaves (BBCH 13-14) until maturity (BBCH 79)	1-3 applications every 15-30 days
Oilseed crops and pulses	Soybean, rapeseed, pea, bean		1.00 %	500-700			
Fruiting vegetables, leaf vegetables, tubers, bulbs, root vegetables							
Industrial crops	Cotton						
Orchards	Stone fruit, pome fruit	~	0.50 %	1000-1500	5-10	From bud bursting to flowering	
	Tropical		0.34 %	1500-2000			1-2 applications every 15-30 days
	Nuts, pomegranate		0.25 %	2000-2500			
Sapling trees		~	0.70 %	700-1000			
Citrus		~	0.20 %	2500-3000			
Olive trees		~	0.25 %	2000-2500			
Grapes	Table grapes Vines	~	0.70 %	700-1000			
Berries	rries		- -100 %	500-700			
Strawberry		~	- 1.00 %	500-700		From first 3-4 leaves (BBCH 13-14) until maturity (BBCH 79)	 1-3 applications every 15-30 days
Foliar application	∆ o Irrigation application			Seed treatment		Furrow treatment	
Optimal dose	Dosage min - max (L/ha)			Treatme	nt volun	ne (L) Treat. Vol (L) min - max	Frequency

If you have any questions, please contact Stoller Europe, or the distributor of Stoller Europe, or your technical advisor, for more specific recommendations.

Observations:

Harbest should be applied from the first phenological states after germination. The product is designed to be applied in foliar sprays, using machinery intended for this purpose and to ensure a sufficient dispersion and drift of droplets to get a homogeneous and efficient application. The product should not be mixed with other liquid fertilisers that contain phosphorous or sulphur in order to avoid insoluble precipitations. However, if mixing is desired, it is advisable to consult your local distributor or representative in advance. These doses are indicative, and it will be the user's responsibility to increase or decrease them according to the crop's characteristics, always following the advice of a specialist technician. In any case, we recommend making a small-scale test before using the product.

RECOMMENDED STORAGE CONDITIONS:

Store the containers between 0 and 35 °C. General conditions for storage: avoid sources of heat, radiation, static electricity and contact with food.

Please also refer to information provided in Material Safety Data Sheet (MSDS).

SAFETY AND ENVIRONMENTAL INFORMATION:

Carefully read the instructions on the label before using the product. To avoid risks to human health and the environment, comply with the instructions for use.

Classification of this product has been carried out in accordance with CLP Regulation (EC) No 1272/2008.



Boric acid (CAS: 10043-35-3)

PRODUCT CLASSIFICATION:

Hazard statements: H319: Causes serious eye irritation. H360FD: May damage fertility. May damage the unborn child. Precautionary statements: P201: Obtain special instructions before use. P264: Wash thoroughly after handling. P280: Wear protective glowes/face protection/protective footwear. P305+P351+P338. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P308+P313: IF exposed or concerned: Get medical advice/attention. P501: Dispose of contents and / or containers in accordance with regulations on hazardous waste or packaging and packaging waste respectively.

This fertiliser contains urea, which can release ammonia and have an impact on air quality. Depending on local conditions, appropriate remediation measures must be taken.

OTHER INFORMATION/ADDITIONAL INFORMATION:

Professional use.

Warning:

Stoller Europe S.L.U. makes all the recommendations based on experience. A lot of uncontrollable factors (preparation of mixtures, climatology, application forms, etc.) may be involved. Therefore, the user will be responsible for any damage that may occur.

ogether we grow.



Manufactured by: STOLLER EUROPE, S.L.U. A Stoller Group Company Elche Parque Empresarial C/ Max Planck, 1. 03203 Elche, Alicante (Spain) (+34) 965 110 522 info@stollereurope.com www.stollereurope.com This document is a reference tool and may only be used as such. The appropriate product label is the one shown on the packaging and printed at the time of purchase.



PHYSICAL FORM: Liquid in solution

NET MASS: 6.7 kg €

51 e

NET VOLUME: LOT NUMBER: