

**COMPOUND INORGANIC MICRONUTRIENT FERTILISER: [PFC 1(C)(II)(b)]
MINERAL MICRONUTRIENT FERTILISER IN SOLUTION****DECLARED CONTENT:**

4.0 % w/wWater-soluble Boron (B) from boron ethanamine
 0.15 % w/wWater-soluble Copper (Cu), from sulphate
 0.015 % w/wWater-soluble Molybdenum (Mo), from sodium
 4.5 % w/wWater-soluble Zinc (Zn), from citrate

Nitrogen (N) or phosphorus pentoxide (P₂O₅) above 0,5 % by mass:

4.0 % w/wNitrogen (N)

LIST OF INGREDIENTS:

Boron ethanamine¹ (CAS: 68425-67-2), Zinc citrate¹(CAS: 546-46-3).

¹Virgin material substances and mixture.

INSTRUCTIONS FOR USE:

SugarMover Zn is a liquid fertilizer that provides boron, copper, molybdenum and zinc, the main nutrients required in the grain and bulb filling process. The boron present in **SugarMover Zn** is an essential element that improves the transport of sugars from the leaves (source) to the grains and bulbs (sinks). Copper, molybdenum and zinc are nutrients that promote photosynthesis and protein synthesis that improve quality, size and uniformity characteristics. Thanks to **Stoller Formulation Technology**, we provide adequate nutrition that naturally intervenes in the physiological processes of plants.

INSTRUCTIONS AND APPLICATION RATES

Crop	☞	☞	☞	☞	☞	Growth stage	☞
Cereals	Wheat, barley, rye, triticale, oats, sorghum, corn, sunflower	✓	0.65 %	300-500	2-3	From seed filling up to maturity (BBCH 49)	1-2 applications every 15-30 days
Fruiting vegetables		✓	0.40 %	700-1000	5-10 3	Color change principle, 30 and 15 days before collection	1-2 applications every 7-15 days
Industrial crops	Cotton	✓	0.70 %	700-1000	5	BBCH 52-61-83 (first appearance of flower buds, beginning of flowering, 30 % cotton bolls open)	1-3 applications every 15-30 days
Oilseed crops and pulses	Soybean, rapeseed, pea, bean	✓	0.30 %	700-1000	2-3	From the appearance of the pods to maturity (BBCH 71-89)	1-2 applications every 15-30 days
Grasses, lucerne		✓	0.65 %	300-500	2-3	10 days before cutting	1 aplicación
Tubers		✓				From tuber filling up to maturity (BBCH 71-89)	
Bulbs		✓	0.30 %	700-1000	2-3	From bulb filling up to maturity (BBCH 71-89)	1-2 applications every 15-20 days
Root vegetables		✓				From root filling up to maturity (BBCH 71-89)	
Orchards	Stone fruit, pome fruit Tropical, avocado, persimmon	✓	0.50 %	1000-1500			
Citrus		✓	0.25 %	2000-2500	5	30 and 15 days before collection	1-2 applications every 7-15 days
Olive trees		✓					
Grapes	Table grapes Vines	✓	1.00 %	500-700			
Berries		✓	0.40 %	700-1000	5-10 3	Before the formation of the inflorescence. Color change principle, 30 and 15 days before collection	2-4 applications every 7-15 days 1-2 applications every 7-15 days

☞ Folar application ☞ Irrigation application ☞ Seed treatment ☞ Furrow treatment
 ☞ Optimal dose ☞ Dosage min - max ☞ Treatment volume (L) ☞ Treat. Vol (L) min - max ☞ Frequency

If you have any questions, please contact Stoller Europe, or the distributor of Stoller Europe, or your technical advisor, for more specific recommendations.

Observations:

The product is compatible with most liquid fertilisers. However, if mixing is desired, it is advisable to consult your local distributor or representative in advance. These doses are indicative, and it will be the user's responsibility to increase or decrease them according to the crop's characteristics, always following the advice of a specialist technician. In any case, we recommend making a small-scale test before using the product. To be used only where there is a recognised need. Do not exceed the application rate.

This document is a reference tool and may only be used as such. The appropriate product label is the one shown on the packaging and printed at the time of purchase.

Manufactured by:
STOLLER EUROPE, S.L.U.
 A Stoller Group Company
 Elche Parque Empresarial
 C/ Max Planck, 1.
 03203 Elche, Alicante (Spain)
 (+34) 965 110 522
 info@stollereurope.com
 www.stollereurope.com

**RECOMMENDED STORAGE CONDITIONS:**

Store the containers between 0 and 35 ° C. General conditions for storage: avoid sources of heat, radiation, static electricity and contact with food. Please also refer to information provided in Material Safety Data Sheet (MSDS).

SAFETY AND ENVIRONMENTAL INFORMATION:

Carefully read the instructions on the label before using the product. To avoid risks to human health and the environment, comply with the instructions for use. Classification of this product has been carried out in accordance with CLP Regulation (EC) No 1272/2008.

**WARNING****PRODUCT CLASSIFICATION:**

Hazard statements: H412: Harmful to aquatic life with long lasting effects. H319: Causes serious eye irritation.
Precautionary statements: P264: Wash thoroughly after handling. P273: Avoid release to the environment. P280: Wear protective gloves/protective clothing/eye protection/protective footwear. P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P337+P313: If eye irritation persists: Get medical advice/attention. P501: Dispose of contents and / or containers in accordance with regulations on hazardous waste or packaging and packaging waste respectively.

OTHER INFORMATION/ADDITIONAL INFORMATION:

Professional use.
 Fertiliser usable in organic plant production according to R(EU) 2018/848.

Warning:

Stoller Europe S.L.U. makes all the recommendations based on experience. A lot of uncontrollable factors (preparation of mixtures, climatology, application forms, etc) may be involved. Therefore, the user will be responsible for any damage that may occur.

PHYSICAL FORM: Liquid in solution
NET MASS: 6.6 kg e
NET VOLUME: 5 L e
LOT NUMBER:

